

TOWARDS A NONSINGULAR UNIVERSE^a

ROBERT H. BRANDENBERGER

Department of Physics

Brown University, Providence, RI 02912, USA

INTRODUCTION

A unified theory of all forces should be nonsingular. In such unified theories, Einstein's theory of general relativity will be a very low curvature effective theory. At larger curvatures, new terms will become important. The question then arises as to whether it is possible to construct an effective action for gravity which leads to a nonsingular theory.

In this work¹⁾ we construct an effective action for gravity in which all homogeneous and isotropic solutions are nonsingular. In particular, there is neither a big bang nor a big crunch. The theory is a higher derivative modification of Einstein's theory constructed in analogy to how the action for point particle motion in special relativity is obtained from Newtonian mechanics. Preliminary work²⁾ indicates that our construction provides a theory in which anisotropies decrease at large curvatures, and in which the inside of a black hole is singularity free.

As is well known, the Penrose-Hawking singularity theorems³⁾ prove that space-time manifolds in general relativity are geodesically incomplete. The key assumptions which are used in the proofs of these theorems is that the action for gravity is the unmodified Einstein action, and that matter satisfies the energy dominance condition $\varepsilon > 0$ and $\varepsilon + 3p \geq 0$, where ε and p are energy density and pressure respectively.

^a Work supported in part by the U.S. Department of Energy under grant DE- AC02-03130, Task A.

In general, the singularity theorems do not provide any information on the nature of the singularity. However, in well known examples such as a collapsing homogeneous and isotropic Universe or the Schwarzschild black hole, at the singularity at least one of the curvature invariants blows up.

Solutions to the “singularity problem” of general relativity such as quantum gravity or string theory all lead to low energy effective actions which include higher derivative terms. As a guide towards constructing an acceptable unified theory we will ask which class of higher derivative effective actions for space-time gives a nonsingular Universe.

Our goal is to construct a theory in which all curvature invariants are bounded and which in addition is geodesically complete. This formidable problem can be reduced substantially by invoking the “Limiting Curvature Hypothesis”⁴⁾, according to which one

- i)* finds a theory in which a small number of specially chosen invariants are explicitly bounded, and
- ii)* when these invariants approach their limiting values, a definite nonsingular solution (namely de Sitter) is taken on.

As a consequence of the limiting curvature hypothesis, automatically all invariants are bounded, and space-time is geodesically complete in its asymptotic regions.

The Limiting Curvature Hypothesis has interesting consequences for Friedman models and for spherically symmetric space-times⁵⁾. A collapsing Universe will not end up in a big crunch, but will approach a contracting de Sitter Universe ($k = 0$) or a de Sitter bounce ($k = 1$) followed by re-expansion (see Fig. 1). For a spherically symmetric vacuum solution, there will be no singularity inside the Schwarzschild horizon; instead, a de Sitter Universe will be reached (see Fig. 1).

Figure 1. Penrose diagrams for a collapsing Universe (left) and for a black hole (right) in Einstein's theory (top) and after implementing the Limiting Curvature Hypothesis (bottom). Wavy lines denote a singularity, the symbols C , DS and E stand for collapsing phase, de Sitter period, and expanding phase respectively, and H denotes the Schwarzschild horizon.

CONSTRUCTION

We shall construct a nonsingular Universe by considering a special class of higher derivative gravity models¹⁾. Higher derivative gravity theories arise as effective actions at high curvatures in many circumstances, *e.g.*, in quantum gravity, when considering quantum fields in an expanding Universe, or in the low energy limit of string theory. It is hence very reasonable to consider such theories.

We construct our special class of models in analogy with how the action

for point particle motion in special relativity can be obtained from Newtonian mechanics⁶⁾. Starting point is the Newtonian action for a point particle with world line $\underline{x}(t)$ and mass m

$$S = m \int dt \frac{1}{2} \dot{x}^2. \quad (1)$$

In this theory, there is no bound on the velocity $\dot{x}$. We construct a new theory by adding a Lagrange multiplier φ which couples to $\dot{x}^2$, the scalar quantity which is to be limited, and give φ a potential $V(\varphi)$. The new action is

$$S = m \int dt \left[\frac{1}{2} \dot{x}^2 + \varphi \dot{x}^2 - V(\varphi) \right]. \quad (2)$$

Provided that $V(\varphi) \sim \varphi$ for $|\varphi| \rightarrow \infty$, the constraint equation (*i.e.*, the variational equation with respect to φ) ensures that $\dot{x}$ is bounded. In order to obtain the correct Newtonian limit for small $\dot{x}$ (*i.e.*, small φ), $V(\varphi)$ must be proportional to φ^2 for $|\varphi| \rightarrow 0$. Up to factors of 2, the simplest potential which satisfies the above asymptotic conditions is

$$V(\varphi) = \frac{2\varphi^2}{1 + 2\varphi}. \quad (3)$$

Eliminating the Lagrange multiplier using the constraint equation and substituting the result into the action, yields the point particle action

$$S = m \int dt \sqrt{1 - \dot{x}^2} \quad (4)$$

in special relativity.

The first step of implementing the Limiting Curvature Hypothesis follows the above procedure. As “old theory” we take Einstein’s action, and from it we construct a new theory in which the Ricci scalar R is limited by introducing a Lagrange multiplier φ_1 with potential $V_1(\varphi_1)$ which couples to R :

$$S = \int d^4x \sqrt{-g} [R + \varphi_1 R + V_1(\varphi_1)] \quad (5)$$

where $V_1(\varphi_1)$ must satisfy the same asymptotic conditions as the multiplier φ in our special relativity example.

As a second step in implementing the Limiting Curvature Hypothesis, we must ensure that as R tends to its limiting values, space-time becomes de Sitter. It is possible to accomplish this by once again using a Lagrange multiplier construction. We find an invariant $I_2(R^{\alpha\beta\gamma\delta})$ with the properties that $I_2 \geq 0$ and $I_2 = 0$ if and only if space-time is de Sitter. We couple a new Lagrange multiplier φ_2 to I_2 and choose its potential $V_2(\varphi_2)$ such that as $|\varphi_2| \rightarrow \infty$ $V_2(\varphi_2) \rightarrow \text{const.}$ Hence, at large $|\varphi_2|$, I_2 is driven to zero and de Sitter space results.

Thus, the new action is

$$S = \int d^4x \sqrt{-g} \left[R + \varphi_1 R - \left(\varphi_2 + \frac{6}{\sqrt{12}} \varphi_1 \right) I_2^{1/2} + V_1(\varphi_1) + V_2(\varphi_2) \right] \quad (6)$$

where the coupling of φ_1 to $I_2^{1/2}$ is introduced to simplify the final equations. For a homogeneous and isotropic metric, the invariant I_2 can be chosen to be

$$I_2 = 4R_{\mu\nu}R^{\mu\nu} - R^2 \quad (7)$$

which is positive for general metrics and vanishes (among homogeneous and isotropic metrics) only for de Sitter spaces.

A simple realization of our scenario uses the potentials

$$\begin{aligned} V_1(\varphi_1) &= 12 H_0^2 \frac{\varphi_1^2}{1 + \varphi_1} \left(1 - \frac{\ln(1 + \varphi_1)}{1 + \varphi_1} \right) \\ V_2(\varphi_2) &= -\sqrt{12} H_0^2 \frac{\varphi_2^2}{1 + \varphi_2^2}. \end{aligned} \quad (8)$$

Applied to a spatially flat collapsing Friedmann-Robertson-Walker metric with scale factor $a(t)$ and Hubble expansion rate $H(t) = \dot{a}(t)/a(t)$, the following variational equations result

$$\begin{aligned} H^2 &= \frac{1}{12} V_1' \\ \dot{H} &= -\frac{1}{\sqrt{12}} V_2' \\ 3(1 - 2\varphi_1)H^2 + \frac{1}{2}(V_1 + V_2) &= \frac{6}{\sqrt{12}} H (\dot{\varphi}_2 + 3H\varphi_2). \end{aligned} \quad (9)$$

From the construction it is already obvious that no singularities arise for trajectories once they reach the asymptotic region $|\varphi_2| \rightarrow \infty$. Equations (9) must be investigated explicitly to show that no singularities arise for finite values of φ_2 , and that trajectories are either periodic about $(\varphi_1, \varphi_2) = (0, 0)$, or else asymptotically approach $|\varphi_2| \rightarrow \infty$.

It has been verified by analytic phase diagram analysis and by numerically solving (9) that indeed no singular points arise in the entire (φ_1, φ_2) plane and that all nonperiodic solutions are asymptotically de Sitter (for more details and figures the reader is referred to Refs. 1 & 7).

It is interesting to include matter in our consideration. Thus, we consider an action

$$S_{\text{full}} = S + S_m \quad (10)$$

where S is the gravitational action (6) and S_m is the matter action for an ideal gas with pressure $p = 0$ (dust) or $p = \frac{1}{3}\rho$ (radiation). In either case, it can be shown⁷⁾ that the asymptotic solutions $|\varphi_2| \rightarrow \infty$ are unchanged. In particular, the system remains singularity free.

This result implies that in our theory, gravity is asymptotically free in the sense that the coupling between matter and gravity becomes negligible at high curvature.

We can also study the variational equations which follow from (6) for a spatially closed homogeneous and isotropic Universe. The only change compared to the spatially flat case is that the de Sitter contraction is replaced by a de Sitter bounce.

SPECULATIONS

If we apply our model (6) to an expanding Universe, we conclude that it has emerged from an initial de Sitter phase, *i.e.*, from an inflationary period. This is not too surprising since many higher derivative gravity theories yield inflation⁸⁾.

However, what is new in our model is that gravity is asymptotically free in the inflationary phase. This might have important consequences for the scalar metric

perturbations produced during inflation⁹⁾. Their amplitude might be suppressed compared to the usual inflationary models. There is the hope of having inflation without the fine tuning problem of density perturbations.

For black holes we conjecture that as the black hole mass shrinks to a mass M_{crit} when the Weyl curvature at the horizon reaches its limiting value, Hawking radiation shuts off. There are no loss of unitarity problems associated with the final stages of black hole evaporation. Finite mass black hole remnants remain, and there is no global charge violation by black holes.

In order for this last conjecture to be reasonable, we must first show that in our theory black hole interiors are nonsingular.

EXTENSIONS

Consider the Schwarzschild metric

$$ds^2 = \left(1 - \frac{2GM}{r}\right) dt^2 - \left(1 - \frac{2GM}{r}\right)^{-1} dr^2 - (dy^2 + dz^2) \quad (11)$$

where y and z are local Cartesian coordinates on S^2 . Inside the horizon, the metric can be approximated by

$$\begin{aligned} ds^2 &\simeq \frac{r}{2GM} dr^2 - \frac{2GM}{r} dt^2 - r^2 (dy^2 + dz^2) \\ &= a^2(\tau) \left[d\tau^2 - e^{-2\beta(\tau)} dx^2 - e^{\beta(\tau)} (dy^2 + dz^2) \right], \end{aligned} \quad (12)$$

where τ is a function of r such that $a(\tau) = \frac{1}{\tau}$ and $r \rightarrow 0$ implies $\tau \rightarrow \infty$. The variable x is a function of t . The above analysis demonstrates that the black hole interior is equivalent to a collapsing Kasner Universe with two scale factors identified.

Hence, if we manage to show that the anisotropy decreases in a collapsing Kasner Universe, we have also shown that the black hole interior will approach de Sitter space, verifying the conjecture of Fig. 1.

Obviously, the invariant I_2 of (7) is insufficient for anisotropic models and for the black hole metric: I_2 vanishes identically for the Schwarzschild metric. However, there is a simple extension:

$$I_2 = 4R_{\mu\nu}R^{\mu\nu} - R^2 + C^2 \quad (13)$$

where $C^2 = C_{\alpha\beta\gamma\delta}C^{\alpha\beta\gamma\delta}$, $C_{\alpha\beta\gamma\delta}$ being the Weyl tensor. For spherical symmetry $C^2 \geq 0$. Hence, forcing $I_2 = 0$ implies both $C^2 = 0$ (vanishing anisotropy) and $4R_{\mu\nu}R^{\mu\nu} - R^2 = 0$ which means that the resulting isotropic space-time is de Sitter²⁾.

To simplify the algebra we can consider a simpler action^{10,2)}:

$$S_A = \int d^4x \sqrt{-g} (R + \varphi C^2 + V(\varphi)) \quad (14)$$

with $V(\varphi) = 1 - \varphi^{-1/2}$ for $|\varphi| \rightarrow \infty$. It is not hard to obtain the variational equations for this problem and to verify that for asymptotic solutions $|\varphi| \rightarrow \infty$ the anisotropy decreases and de Sitter space is approached.

Since these results should carry over to our system of (6) and (13), we conclude that in our model, all homogeneous solutions will approach de Sitter space, and that the black hole interior will be nonsingular.

CONCLUSIONS AND ACKNOWLEDGEMENTS

We have constructed an effective action for gravity, based on a higher derivative modification of Einstein's theory of general relativity, in which all homogeneous solutions are nonsingular. They are either periodic about Minkowski space-time, or else they asymptotically approach de Sitter space. We have given some preliminary evidence that in our model also singularities inside of black holes can be avoided.

The results and ideas presented in this lecture are based on key ideas by and joint work with V. Mukhanov. I am grateful to him and to my collaborators M. Mohazzab, A. Sornborger and M. Trodden.

REFERENCES

1. V. Mukhanov and R. Brandenberger, *Phys. Rev. Lett.* **68**, 1969 (1992).
2. R. Brandenberger, M. Mohazzab, V. Mukhanov, A. Sornberger and M. Trodden, in preparation (1993).
3. R. Penrose, *Phys. Rev. Lett.* **14**, 57 (1965);
S. Hawking, *Proc. R. Soc. London* **A300**, 182 (1967).
4. M. Markov, *Pis'ma Zh. Eksp. Teor. Fiz.* **36**, 214 (1982); **46**, 342 (1987);
V. Ginsburg, V. Mukhanov and V. Frolov, *Zh. Eksp. Teor. Fiz.* **94**, 1 (1988).
5. V. Frolov, M. Markov and V. Mukhanov, *Phys. Lett.* **B216**, 272 (1989);
Phys. Rev. **D41**, 383 (1990).
6. B. Altshuler, *Class. Quant. Grav.* **7**, 189 (1990).
7. R. Brandenberger, V. Mukhanov and S. Sornborger, "A Cosmological Theory Without Singularities" Brown preprint BROWN-HET-891 (1993).
8. A. Starobinskii, *Phys. Lett.* **B91**, 99 (1980).
9. V. Mukhanov, H. Feldman and R. Brandenberger, *Phys. Rep.* **215**, 203 (1992).
10. B. Altshuler, in Proc. of the 1st Int. A.D. Sakharov Conference, Moscow, May 1991, to be published in the proceedings (Nova Science Publ., Inc., New York, 1992)